

ON NORMAL FERMI-WALKER DERIVATIVE FOR FOCAL CURVES WITH RIBBON FRAME

MUSTAFA YENEROĞLU AND VEDAT ASIL

ABSTRACT. In this paper, we study a new construction of curves by Normal Fermi-Walker parallelism and derivative with Ribbon frame. Finally, we give Normal Fermi-Walker parallelism and derivative for a focal curve according to Ribbon frame.

1. INTRODUCTION

Fermi-Walker parallelism and derivative are extremely vital for various useful approaches. Fermi-Walker parallelism and derivative are very important in many practical applications. Fermi Walker derivative in the Lorentzian manifolds is a generalization of covariant differentiation. The derivatives of the spacelike unit vectors in general relativity is taken with respect to timelike unit vectors. Additionally, this derivative is used to define non-inertial but non-rotation frames. In the special case of inertial frames, the dervative degrade covariant derivative. Fermi Walker derivative along any curve preserves the orthogonality of the tangent space parellel. A large class of reference frames in space-time experience inertial forces and therefore constitute noninertial frames. For instance, the frame adapted to an observer “at rest” on the surface of the (rotating) Earth is noninertial. Observers that follow arbitrary timelike trajectories in space-time will regard as natural a reference frame in which they are at rest, and their spatial axes do not rotate. Fermi–Walker transported frames are important in several investigations. All inner products are invariant under this transport [13]. A frame that undergoes linear and rotational acceleration may be described by the Frenet– Serret frame. In [10], authors considered tetrad fields as reference frames adapted to observers that move along arbitrary timelike trajectories in space-time. By means of a local Lorentz transformation, they transformed these frames into Fermi–Walker transported frames. Moreover, they presented a simple prescription for the construction of Fermi–Walker transported frames out of an arbitrary set of tetrad fields. Recently, Karakuş, Yaylı and Körpınar have made several studies related to Fermi-Walker derivation [6,7].

On the other hand, for any unit tangent vector of the curve , the focal curve is defined as the centers of the osculating spheres. R.U. Vargas examined the properties of the focal curve [12]. G. Öztürk and K. Arslan gave some results about

1991 *Mathematics Subject Classification*. Primary 53A04.

Key words and phrases. Ribbon frame, Normal Fermi Walker derivative-parallelism, Focal curve.

the focal curve in Euclidean n -space [11-14]. N. Gürses, Ö. Bektaş and S. Yüce studied the Quaternionic Focal Curves [17]. Furthermore, authors gave differential applications of the curve and surfaces [5-15,16,15-27].

Ribbon theory may be applicable in DNA and protein folding studies [7,8]. In addition to, the differential geometry of ribbons have been considered in connection with statical models for models for polymers [9]. Bohr and Markvorsen also studied ribbon frame [4].

The main goal this article is analyzed normal Fermi-Walker derivations along the focal curve with ribbon frame.

2. PRELIMINARIES

To construct any ribbon – and hence its center curve – we will use two continuous functions, $w(s)$ and $\theta(s)$ defined in the interval $s \in [0, L]$, where L is the intrinsic length of the ribbon under construction. Throughout we assume that $\theta(s) \in]0, \pi[$ for all s so that in particular $\sin(\theta(s)) > 0$ for all s . In our construction a key role is played by a unit vector field $\mathbf{a}(s)$, which is a field tangent to the ribbon and defined by having the angle $\theta(s)$ to the center curve of the ribbon. In fact, $\mathbf{a}(s)$ will be the direction field for the Darboux vector $\mathbf{D}(s)$ with the generating function $w(s)$ as a multiplying factor, i.e. $\mathbf{D}(s) = w(s)\mathbf{a}(s)$, [4].

We let $\{e(s), f(s), g(s)\}$ denote the unique orthonormal triple of vector solutions to the following differential system:

$$\begin{aligned} e'(s) &= w(s)a(s) \times e(s), \\ f'(s) &= w(s)a(s) \times f(s), \\ g'(s) &= w(s)a(s) \times g(s), \end{aligned}$$

where the unit vector field $\mathbf{a}(s)$ is defined in terms of $e(s)$ and $g(s)$ as follows:

$$\mathbf{a}(s) = \cos(\theta(s))e(s) + \sin(\theta(s))g(s),$$

and where – for uniqueness purposes – we also apply the following arbitrary initial conditions referring to a given fixed coordinate system and basis in \mathbb{R}^3

$$\{e(0), f(0), g(0)\} = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}.$$

The above system can be written explicitly as follows:

$$\begin{aligned} e'(s) &= w(s) \sin(\theta(s))f(s), \\ f'(s) &= -w(s) \sin(\theta(s))e(s) + w(s) \cos(\theta(s))g(s), \\ g'(s) &= -w(s) \cos(\theta(s))f(s), \end{aligned}$$

Definition 2.1. X is any vector field and $\alpha(s)$ is unit speed any curve in space.

$$\frac{\tilde{D}X}{\tilde{D}s} = \frac{dX}{ds} - \langle N, X \rangle N' + \langle N', X \rangle N$$

defined as $\frac{\tilde{D}X}{\tilde{D}s}$ derivative is called Normal Fermi-Walker derivative. Here $N = \frac{T'}{\|T'\|}$ is a normal vector Frenet frame [25].

Definition 2.2. X is any vector field along the $\alpha(s)$ space curve. If the Normal Fermi-Walker derivative of the vector field X

$$\frac{\tilde{D}X}{\tilde{D}s} = 0$$

the vector field X along the curve is called normal Fermi-Walker parallel [25].

3. Ribbon Frame and Normal Fermi-Walker Derivative

According to ribbon frame, normal Fermi Walker derivative of the vector field X is defined as follows:

$$\frac{\tilde{D}X}{\tilde{D}s} = \frac{dX}{ds} - \langle f, X \rangle f' + \langle f', X \rangle f$$

Theorem 3.1. Let $\{e, f, g\}$ and X be ribbon frame and any vector field along the $\alpha(s)$ space curve, respectively . Normal Fermi-Walker derivative can be expressed along the curve on ribbon frame as follows:

$$\frac{\tilde{D}X}{\tilde{D}s} = \frac{dX}{ds} + w(s) \sin \theta(s)(X \wedge g) + w(s) \cos \theta(s)(X \wedge e)$$

Proof:

$$\begin{aligned} \frac{\tilde{D}X}{\tilde{D}s} &= \frac{dX}{ds} - \langle f, X \rangle (-w(s) \sin \theta(s)e + w(s) \cos \theta(s)g) + \langle (-w(s) \sin \theta(s)e + w(s) \cos \theta(s)g), X \rangle f, \\ &= \frac{dX}{ds} + w(s) \sin \theta(s) \langle f, X \rangle e - w(s) \cos \theta(s) \langle f, X \rangle g - w(s) \sin \theta(s) \langle e, X \rangle f + w(s) \cos \theta(s) \langle g, X \rangle f, \\ &= \frac{dX}{ds} + w(s) \sin \theta(s) [\langle f, X \rangle e - \langle e, X \rangle f] + w(s) \cos \theta(s) [\langle g, X \rangle f - \langle f, X \rangle g] \\ &= \frac{dX}{ds} + w(s) \sin \theta(s)(X \wedge g) + w(s) \cos \theta(s)(X \wedge e) \end{aligned}$$

Theorem 3.2. Let $\alpha(s)$ be a unit-speed curve. Then $X = \lambda_1 e + \lambda_2 f + \lambda_3 g$ vector field according to ribbon frame along the curve is parallel to normal Fermi-Walker terms if and only if:

$$\begin{aligned} \lambda_1 &= \text{cons.}, \\ \lambda_2 &= \text{cons.}, \\ \lambda_3 &= \text{cons.}, \end{aligned}$$

where $\lambda_1, \lambda_2, \lambda_3$ are continuously differentiable functions according to real parameter s .

Proof. \implies :The vector field X is parallel to the normal Fermi-Walker terms. Then

$$\begin{aligned} \frac{\tilde{D}X}{\tilde{D}s} &= \frac{dX}{ds} + w(s) \sin \theta(s)(X \wedge g) + w(s) \cos \theta(s)(X \wedge e) \\ \frac{\tilde{D}X}{\tilde{D}s} &= \frac{d\lambda_1}{ds} e + \frac{d\lambda_2}{ds} f + \frac{d\lambda_3}{ds} g \end{aligned}$$

is obtained. X is parallel to the Fermi-Walker terms and $\frac{\tilde{D}X}{\tilde{D}s} = 0$ so,

$$\begin{aligned} \frac{d\lambda_1}{ds} &= 0 \\ \frac{d\lambda_2}{ds} &= 0 \\ \frac{d\lambda_3}{ds} &= 0 \end{aligned}$$

is obtained. This the solution of the equation system is given as follows:

$$\begin{aligned} \lambda_1 &= \text{cons.}, \\ \lambda_2 &= \text{cons.}, \\ \lambda_3 &= \text{cons.}, \end{aligned}$$

\Leftarrow : $X = \lambda_1 e + \lambda_2 f + \lambda_3 g$ vector field and

$$\begin{aligned} \lambda_1 &= \text{cons.}, \\ \lambda_2 &= \text{cons.}, \\ \lambda_3 &= \text{cons.}, \end{aligned}$$

and from theorem 3.1,

$$\frac{\tilde{D}X}{\tilde{D}s} = 0$$

is obtained.

4. Focal Curve According to Ribbon Frame and Normal Fermi-Walker Derivative

Denoting the focal curve by $C_\alpha^{\mathcal{R}}$ according to ribbon frame, we can write

$$C_\alpha^{\mathcal{R}}(s) = (\alpha + c_1 f + c_2 g)(s),$$

where the coefficients $c_1 = \frac{1}{w(s) \sin(\theta(s))}$, $c_2 = \frac{c'_1}{w(s) \cos(\theta(s))}$ are smooth functions of the parameter of the curve α , called the first and second focal curvatures of α , respectively. Frenet vectors of the focal curve $C_\alpha^{\mathcal{R}}$ according to ribbon frame are given as follows:

$$\begin{aligned} T_F &= (c'_2 + \cot \theta(s))g \\ N_F &= (c'_2 + \cot \theta(s))'g - (c'_2 + \cot \theta(s))w(s) \cos \theta(s)f \\ B_F &= (c'_2 + \cot \theta(s))^2 w(s) \cos \theta(s)e \end{aligned}$$

and

$$\begin{aligned} T'_F &= \kappa_F N_F \\ N'_F &= -\kappa_F T_F + \tau_F B_F \\ B'_F &= -\tau_F N_F \end{aligned}$$

Theorem 4.1. Let $\{T_F, N_F, B_F\}$ and X be Frenet frame and any vector field along the $C_\alpha^{\mathcal{R}}$ focal curve, respectively. Normal Fermi-Walker derivative can be

expressed along the curve on this frame as follows:

$$\frac{\tilde{D}X}{\tilde{D}s} = \frac{dX}{ds} - \kappa_F(c'_2 + \cot \theta(s))^2 w(s) \cos \theta(s) X \wedge e + \tau_F(c'_2 + \cot \theta(s)) X \wedge g$$

Proof.

$$\begin{aligned} \frac{\tilde{D}X}{\tilde{D}s} &= \frac{dX}{ds} - \langle N_F, X \rangle N'_F + \langle N'_F, X \rangle N_F, \\ \frac{\tilde{D}X}{\tilde{D}s} &= \frac{dX}{ds} - \langle N_F, X \rangle (-\kappa_F T_F + \tau_F B_F) + \langle -\kappa_F T_F + \tau_F B_F, X \rangle N_F, \\ \frac{\tilde{D}X}{\tilde{D}s} &= \frac{dX}{ds} + \kappa_F X \wedge B_F + \tau_F X \wedge T_F \\ \frac{\tilde{D}X}{\tilde{D}s} &= \frac{dX}{ds} - \kappa_F(c'_2 + \cot \theta(s))^2 w(s) \cos \theta(s) X \wedge e + \tau_F(c'_2 + \cot \theta(s)) X \wedge g \end{aligned}$$

Theorem 4.2. Let C_α^R be a unit-speed focal curve. Then $X = \lambda_1 T_F + \lambda_2 N_F + \lambda_3 B_F$ vector field according to Frenet frame along the focal curve is parallel to normal Fermi-Walker terms if and only if:

$$\begin{aligned} \lambda_1 &= \text{cons.}, \\ \lambda_2 &= \text{cons.}, \\ \lambda_3 &= \text{cons.}, \end{aligned}$$

where $\lambda_1, \lambda_2, \lambda_3$ are continuously differentiable functions according to real parameter s .

Proof. \implies : The vector field X is parallel to the normal Fermi-Walker terms. Then

$$\frac{\tilde{D}X}{\tilde{D}s} = \frac{d\lambda_1}{ds} T_F + \frac{d\lambda_2}{ds} N_F + \frac{d\lambda_3}{ds} B_F$$

is obtained. X is parallel to the Fermi-Walker terms and $\frac{\tilde{D}X}{\tilde{D}s} = 0$ so,

$$\begin{aligned} \frac{d\lambda_1}{ds} &= 0 \\ \frac{d\lambda_2}{ds} &= 0 \\ \frac{d\lambda_3}{ds} &= 0 \end{aligned}$$

is obtained. This the solution of the equation system is given as follows:

$$\begin{aligned} \lambda_1 &= \text{cons.}, \\ \lambda_2 &= \text{cons.}, \\ \lambda_3 &= \text{cons.}, \end{aligned}$$

\Leftarrow : $X = \lambda_1 T_F + \lambda_2 N_F + \lambda_3 B_F$ vector field

$$\lambda_1 = \text{cons.},$$

$$\lambda_2 = \text{cons.},$$

$$\lambda_3 = \text{cons.},$$

and from theorem 4.1,

$$\frac{\tilde{D}X}{\tilde{D}s} = 0$$

is obtained.

REFERENCES

- [1] E. Fermi, *Atti Accad. Naz. Lincei Cl. Sci. Fiz. Mat. Nat.* 31 (1922) 184–306.
- [2] A. Gray: *Modern Differential Geometry of Curves and Surfaces with Mathematica*. CRC Press (1998).
- [3] J.W. Maluf and F. F. Faria, On the construction of Fermi-Walker transported frames, *Ann. Phys. (Berlin)* 17 (5) (2008), 326 – 335
- [4] J. Bohr and S. Markvorsen, *Ribbon Crystals*, *Plos one*, 8 (10) (2013).
- [5] T. Körpınar, New Characterization for Minimizing Energy of Biharmonic Particles in Heisenberg Spacetime, *Int J Phys.*53 (2014) 3208-3218
- [6] F. Karakuş and Y. Yaylı, On the Fermi-Walker Derivative and Non-Rotating Frame, *World Sci.* 9(8) (2012).
- [7] G.S. Chirikjian, Framed curves and knotted DNA, *Biochem Soc. Trans.* 41 (2013), 635-638
- [8] S. Hu, M. Lundgren, A.J. Niemi, Discrete frenet frame, inflection point solutions, and curve visualization with applications to folded proteins, *Phys. Rev. E.* 83 (2011).
- [9] S.M Rappaport, Y. Rabin, Differential geometry of polymer models: worm-like chains, ribbons and fourier knots, *J Phys A: Math Theory* 40 (2007).
- [10] J.W. Maluf and F.F. Faria, On the construction of Fermi-Walker transported frames, *Ann. der Phys.*, 17, 326 (2008).
- [11] P. Alegre, K. Arslan, A. Carriazo, C. Murathan and G. Öztürk, Some pecial types of the developable ruled surface, *Hacettepe J. Math. and Statis.*, 39(3), (2010).
- [12] R. Uribe-Vargas, On vertices, focal curvatures and differential geometry of space curves, *Bull. Brazilian Math. Soc.* 36 (3) (2005), 285–307.
- [13] D. Bini and R.T. Jantzen, Circular holonomy, clock effects and gravitoelectromagnetism: still going around in circles after all these years..., *Proc. of the 9th ICRA Workshop on Fermi and Astrophysics*, 2001, eds. R.Ruffini and C.Sigismondi (World Scientific,2002)
- [14] G. Öztürk and K. Arslan, On focal curves in Euclidean n-space \mathbb{R}^n , *Novi Sad Journal of Mathematics*, 46 (1), 35-44 (2016).
- [15] T. Körpınar, B-tubular surfaces in Lorentzian Heisenberg Group H3, *Acta Scientiarum. Technology* 37(1) (2015), 63–69
- [16] T. Körpınar, S. Bas, On Characterization Of B- Focal Curves In E^3 , *Bol. Soc. Paran. Mat.*, 31 (1) (2013), 175-178.
- [17] N. Gürses, Ö. Bektaş and S. Yüce, On the Quaternionic Focal Curves, *Süleyman Demirel University Journal of Natural and Applied Sciences*, 21(2) (2017), 357-366.
- [18] T. Körpınar, On the Fermi-Walker Derivative for Inextensible Flows, *Zeitschrift für Naturforschung A.* 70 (7) (2015), 477–482
- [19] G. Altay, H. Öztekin, Translation Surfaces Generated by Mannheim Curves in Three Dimensional Euclidean Space , *Gen. Math. Notes*, 26 (1),2015, 28-34.
- [20] V. Asil, T. Körpınar, Dual focal curves of dual spacelike curves in the Dual Lorentzian D13 *Palestine Journal of Mathematics* , 2 (1) (2013), 1-3.
- [21] T. Körpınar, E. Turhan and S. Baş: Characterizing Of Dual Focal Curves In D3, *Bol. Soc. Paran. Mat.*, 31 (2) (2013), 77–82.

- [22] T. Körpınar, E. Turhan: Construction of B-Focal Curves of Spacelike Biharmonic B-slant Helices According to Bishop Frame in $E(1,1)$, AAPP, DOI:10.1478/AAPP.901A4
- [23] V. Asil, T. Körpınar, S. Bař: Construction of dual focal curves in $D3$, Advanced Modeling and Optimization, 14 (2) (2012), 293-296.
- [24] T. Körpınar, E. Turhan : On characterization of dual focal curves of spacelike biharmonic curves with timelike binormal in the dual lorentzian $D31$, Stud. Univ. Babeř-Bolyai Math. 57 (3) (2012), 421–426.
- [25] Ö. Keskin and Y. Yaylı, Normal Fermi Walker Derivative, Mathematical Sciences and Applications E-Notes, 5(1), 1-8, (2017).
- [26] T. Körpınar: A New Method for Inextensible Flows of Timelike Curves in Minkowski Space-Time E_1^4 , International Journal of Partial Differential Equations, Volume 2014, Article ID 517070, 7 pages
- [27] T. Körpınar, Bianchi Type-I Cosmological Models for Inextensible Flows of Biharmonic Particles by Using Curvature Tensor Field in Spacetime, Int J Theor Phys 54 (2015), 1762–1770

FIRAT UNIVERSITY, DEPARTMENT OF MATHEMATICS, 23119, ELAZIĞ, TURKEY
E-mail address: mustafayenerglu@gmail.com, vasil@firat.edu.tr